

Evolution of non-linear cosmological perturbations

David Langlois^{1,2}, Filippo Vernizzi³

¹*APC (Astroparticules et Cosmologie), UMR 7164 (CNRS, Université Paris 7, CEA, Observatoire de Paris), 11 Place Marcelin Berthelot, F-75005 Paris, France*

²*Institut d'Astrophysique de Paris, 98bis Boulevard Arago, 75014 Paris, France, and*

³*Helsinki Institute of Physics, P.O. Box 64, FIN-00014, University of Helsinki, Finland*

We define fully non-perturbative generalizations of the uniform density and comoving curvature perturbations, which are known, in the linear theory, to be conserved on sufficiently large scales for adiabatic perturbations. Our non-linear generalizations are defined geometrically, independently of any coordinate system. We give the equations governing their evolution on all scales. Also, in order to make contact with previous works on first and second order perturbations, we introduce a coordinate system and show that previous results can be recovered, on large scales, in a remarkably simple way, after restricting our definitions to first and second orders in a perturbative expansion.

The relativistic theory of cosmological perturbations is a cornerstone in our understanding of the early universe as it is indispensable in relating the scenarios of the early universe, such as inflation, to cosmological data such as the Cosmic Microwave Background (CMB) anisotropies. This is the reason why the theory of *linear* cosmological perturbations has been developed to a high degree of sophistication during the last twenty-five years [1, 2, 3].

With the increase of precision of the CMB data, the study of relativistic cosmological perturbations *beyond linear order* is becoming a topical subject, especially to study primordial non-Gaussianity. This is why much effort has been devoted recently to the investigation of second-order perturbations, mainly resorting to an expansion, up to second order, of Einstein's equations [4, 5]. This formalism is rather difficult, due to the complexity of Einstein's equations, and the identification of gauge invariant quantities is not straightforward [6, 7].

Other recent approaches of non-linear perturbations [8, 9, 10] are based on the long wavelength approximation, which at lowest order is related to the so called separate universe picture that represents our universe, on scales larger than the Hubble radius, as juxtaposed Friedmann-Lemaître-Robertson-Walker (FLRW) universes with slightly different scale factors.

In the present work, we propose a different approach based on a purely geometrical description of the perturbations. Our approach does not rely on any perturbative expansion and is thus not limited to second-order perturbations. It does not require any approximation and covers both large and small scales. Furthermore, as we shall see, it can be related rather easily to the formulations mentioned above. Our approach is directly inspired by the so called covariant formalism for cosmological perturbations introduced by Ellis and Bruni [11]. At linear order, the covariant formalism is *computationally* equivalent to the much more used coordinate approach. However, as we show here, it is tremendously more powerful when one goes beyond linear order.

In particular, we define geometrically the generalizations of quantities widely used in the linear theory be-

cause they are conserved on large scales. We then give the full non-linear equations that govern these quantities. To emphasize the efficiency of our approach with respect to the more traditional coordinate-based approach, we show how second order gauge invariant quantities, recently derived in the literature, can be recovered, on large scales, from our geometric quantities via a straightforward derivation.

Covariant approach – We consider a spacetime, with metric g_{ab} , filled with a single perfect fluid characterized by a four-velocity $u^a = dx^a/d\tau$ ($u_a u^a = -1$), where τ is the proper time along the flow lines, proper energy density ρ and pressure P . The corresponding energy-momentum tensor is $T^a_b = (\rho + P)u^a u_b + P g^a_b$. We also introduce the volume expansion $\Theta \equiv \nabla_a u^a$, which reduces to the Hubble parameter H , $\Theta = 3H$, in a FLRW spacetime. The integration of Θ along the fluid world lines, with respect to the proper time τ ,

$$\alpha \equiv \frac{1}{3} \int d\tau \Theta, \quad (\Theta = 3\dot{\alpha} \equiv 3u^a \nabla_a \alpha), \quad (1)$$

can be used to define, for each observer comoving with the fluid, a local scale factor e^α . Note that α is defined up to an integration constant for *each* fluid world line, which we can set to zero on some reference hypersurface.

As shown by Ellis and Bruni [11], it is possible to define, in a geometrical way, quantities that can be interpreted as perturbations with respect to the FLRW configuration, by introducing spatially projected gradients. The spatial projection tensor, orthogonal to the fluid velocity u^a , is defined by $h^a_b = g^a_b + u^a u_b$. Defining the spatial gradient $D_a \equiv h^b_a \nabla_b$, we consider the quantities

$$X_a \equiv D_a \rho, \quad Y_a \equiv D_a P, \quad Z_a \equiv D_a \Theta, \quad W_a \equiv D_a \alpha,$$

which automatically vanish in a strictly FLRW spacetime: in this sense we call them *perturbations*. However, they are fully non-perturbative quantities. Note that the last quantity, W_a , which as far as we know has not been introduced previously in the covariant formalism, plays a crucial rôle in our approach, where we replace the familiar spatial curvature perturbation by the perturbation of

the integrated expansion α , i.e., of the local number of e-folds, in accordance with the separate universe picture [12, 13].

Evolution of perturbations – Starting from the above definitions, we now derive the fundamental evolution equation for these non-linear perturbations. As in [13] in the context of the linear theory, our starting point here will be the energy-momentum conservation,

$$\nabla_a T^a_b = 0. \quad (2)$$

The projection along u^a of Equation (2) yields

$$\dot{\rho} + \Theta(\rho + P) = 0,$$

where a dot represents a derivative with respect to τ , i.e., stands for $u^a \nabla_a$. If one takes the projected gradient of this expression one gets, after some manipulations, the relation

$$\dot{X}_a + (\rho + P) Z_a + \Theta(X_a + Y_a) + (h_a^c \nabla_c u^b - \dot{h}_a^b) \nabla_b \rho = 0. \quad (3)$$

We then rewrite Z_a as

$$Z_a = 3\dot{W}_a + 3(h_a^c \nabla_c u^b - \dot{h}_a^b) \nabla_b \alpha. \quad (4)$$

This suggests introducing the covector

$$\zeta_a \equiv W_a + \frac{X_a}{3(\rho + P)} = D_a \alpha - \frac{\dot{\alpha}}{\dot{\rho}} D_a \rho, \quad (5)$$

which represents our *non-perturbative generalization of the curvature perturbation on uniform density hypersurfaces* or, more exactly, of its “spatial” gradient. Interestingly, the same linear combination has been introduced in the context of the long wavelength approximation, within a specific coordinate system, in [8]. Here, ζ_a is defined geometrically and in the most general context.

The spatial projection of Eq. (3), via h_a^b , then yields

$$e^{-\alpha} h_a^b (e^\alpha \zeta_b)^\cdot = -\frac{\Theta}{3(\rho + P)} \Gamma_a - \zeta^b (\sigma_{ba} + \omega_{ba}), \quad (6)$$

where we have used the familiar decomposition (see e.g. [14]) $\nabla_b u_a = \sigma_{ab} + \omega_{ab} + \frac{1}{3}\Theta h_{ab} - \dot{u}_a u_b$ with the (symmetric) shear tensor σ_{ab} , and the (antisymmetric) vorticity tensor ω_{ab} ; on the right hand side, the quantity

$$\Gamma_a \equiv Y_a - c_s^2 X_a \equiv D_a P - \frac{\dot{P}}{\dot{\rho}} D_a \rho \quad (7)$$

represents the non-perturbative generalization of the nonadiabatic pressure and vanishes for purely adiabatic perturbations, i.e., when the pressure P is solely a function of the density ρ . Note also that, for both ζ_a and Γ_a , one can replace the projected derivatives by ordinary partial derivatives respectively in Eqs. (5) and (7).

Equation (6) is one of the main results of this work. It gives the fully non-linear evolution equation for the

variable ζ_a and corresponds to the non-perturbative generalization of the familiar linear conservation law for adiabatic perturbations ($\Gamma_a = 0$) on large scales, similar to the results recently obtained in the long wavelength limit [8].

Introducing coordinates – To relate our approach to the more familiar coordinate-based perturbative approach, we now introduce a coordinate system, in which the perturbed metric can be written as $ds^2 = a^2[-(1+2A)d\eta^2 + 2B_i dx^i d\eta + (\gamma_{ij} + \delta\gamma_{ij}) dx^i dx^j]$ (see [15]).

In the following, we decompose any quantity χ in the form $\chi(\eta, x^i) = \bar{\chi}(\eta) + \delta\chi_1(\eta, x^i) + \frac{1}{2}\delta\chi_2(\eta, x^i)$, where $\delta\chi_1$ and $\delta\chi_2$ represent, respectively, the first and second order perturbations. Let us write down explicitly the components of our covector ζ_a in this coordinate system. The zeroth order vanishes and, at *first order*, the spatial components are simply

$$\zeta_i^{(1)} = \partial_i \zeta_1, \quad \zeta_1 \equiv \delta\alpha_1 - \frac{\bar{\alpha}'}{\rho'} \delta\rho_1, \quad (8)$$

where a prime denotes the partial derivative with respect to the time coordinate η . To compute the component ζ_0 , it is useful to note that for any function χ , one can write $D_0 \chi = u_0 u^i \partial_i \chi - u^i u_i \partial_0 \chi$, where we have used the normalization of u^a . Since u^i is first order, this implies that $\zeta_0^{(1)} = 0$ and $\zeta_0^{(2)} = u^i \zeta_i^{(1)}$.

Expanding $\zeta_i = \partial_i \alpha - (\dot{\alpha}/\dot{\rho}) \partial_i \rho$ at *second order*, one finds, after some simple manipulations,

$$\zeta_i^{(2)} = \partial_i \zeta_2 + \frac{2}{\rho'} \delta\rho_1 \partial_i \zeta_1', \quad (9)$$

$$\zeta_2 \equiv \delta\alpha_2 - \frac{\bar{\alpha}'}{\rho'} \delta\rho_2 - \frac{2}{\rho'} \delta\alpha_1' \delta\rho_1 + 2 \frac{\bar{\alpha}'}{\rho'^2} \delta\rho_1 \delta\rho_1' + \frac{1}{\rho'} \left(\frac{\bar{\alpha}'}{\rho'} \right)' \delta\rho_1^2. \quad (10)$$

Note that the conservation of $\zeta_i^{(1)}$ and $\zeta_i^{(2)}$ is equivalent to the conservation of ζ_1 and ζ_2 .

We now relate α to the metric perturbations, considering for simplicity only large scales. Neglecting gradients as well as first order vector and tensor perturbations, and writing $\psi \equiv -\gamma^{ij} \delta\gamma_{ij}/3$, one can use the equality $3\dot{\alpha} = \nabla_a u^a$ to derive, *up to second order*, the relation

$$\alpha \simeq \ln a - \psi - \psi^2. \quad (11)$$

Thus, from Eq. (10), we can relate ζ_2 to the quantity defined by Malik and Wands in [6],

$$\zeta_2 \simeq \zeta_{2(\text{MW})} - 2\zeta_{1(\text{MW})}^2.$$

The expansion of the nonadiabatic term $\Gamma_a = \partial_a P - (\dot{P}/\dot{\rho}) \partial_a \rho$, which can be read from the expansion of ζ_a by substituting P to α , enables us to write explicitly the conservation equation (6) at first and second orders. If one ignores the second term on the right hand side of

Eq. (6), one finds, at first and second order, respectively,

$$\begin{aligned}\zeta'_1 &\simeq -\frac{\mathcal{H}}{\bar{\rho} + \bar{P}}\Gamma_1, \\ \zeta'_2 &\simeq -\frac{\mathcal{H}}{\bar{\rho} + \bar{P}}\Gamma_2 - \frac{2}{\bar{\rho} + \bar{P}}\Gamma_1\zeta'_1,\end{aligned}$$

where the $\Gamma_{1,2}$ are defined as the $\zeta_{1,2}$, and $\mathcal{H} \equiv a'/a$. One thus recovers the results of [6] very easily. Moreover, Eq. (6) generalizes them to any order in the perturbation theory and to all scales.

Comoving “curvature” perturbation – In addition to the uniform energy density curvature perturbation, generalized by our ζ_a , another useful quantity in the linear theory is the curvature perturbation on comoving hypersurfaces, usually denoted by $\mathcal{R}$. In our formulation, since the perturbed integrated expansion α replaces the usual spatial curvature and since our spatial gradients are defined with respect to the comoving observers, the generalization of $\mathcal{R}$ is simply

$$\mathcal{R}_a \equiv -D_a\alpha = -W_a. \quad (12)$$

In the coordinate system introduced earlier, the spatial components are

$$\mathcal{R}_i = -\partial_i\alpha - \dot{\alpha}a(v_i + B_i),$$

with $v^i \equiv au^i$. At first order, $\delta\alpha \simeq -\psi$, and one recognizes the familiar comoving curvature perturbation of the linear theory.

It is relatively easy to derive the evolution equation for $\mathcal{R}_a$. Equation (4) can be seen as an equation for $\dot{\mathcal{R}}_a = -\dot{W}_a$. Projecting orthogonally to u^a , we find

$$h_a^b \dot{\mathcal{R}}_b + \frac{1}{3}\Theta\mathcal{R}_a = -\mathcal{R}^b(\sigma_{ba} + \omega_{ba}) - h_a^b \dot{u}_b \dot{\alpha} - \frac{1}{3}Z_a. \quad (13)$$

The term involving Z_a on the right hand side can be rewritten by using the shear constraint equation [see Eq. (4.17) of [14]],

$$\frac{2}{3}Z_a - h_a^b \nabla_c(\sigma_b^c + \omega_b^c) + (\sigma_a^b + \omega_a^b)\dot{u}_b = q_a = 0, \quad (14)$$

where q_a is the energy flux in the energy-momentum tensor, which, in our case, vanishes. Moreover, one can replace $\dot{u}_b$ using the Euler equation, $\dot{u}_a = -Y_a/(\rho + P)$. On rewriting Y_a as $Y_a = \Gamma_a + c_s^2 X_a$, and combining all this into Eq. (13), we finally obtain

$$\begin{aligned}e^{-\alpha}h_a^b(e^\alpha\mathcal{R}_b)^\cdot &= \left[\frac{\Theta\delta_a^b}{3(\rho + P)} - \frac{\sigma_a^b + \omega_a^b}{2(\rho + P)} \right] (\Gamma_b + c_s^2 X_b) \\ &\quad - \mathcal{R}^b(\sigma_{ba} + \omega_{ba}) - \frac{1}{2}h_a^b \nabla_c(\sigma_b^c + \omega_b^c)\end{aligned} \quad (15)$$

This is the fully non-linear evolution equation for $\mathcal{R}_a$. As one can see, it is slightly more complicated than the evolution equation for ζ_a . However, when σ_{ab} and ω_{ab}

are negligible, one can recover the familiar conservation law for adiabatic perturbations by noting that X_a , which generalizes the comoving energy density perturbation of the linear theory [2], can be shown to be negligible on large scales. Since $\zeta_a + \mathcal{R}_a = -(\dot{\alpha}/\dot{\rho})X_a$, one then finds $\zeta_a \simeq -\mathcal{R}_a$ non-perturbatively.

Note that $\mathcal{R}_a$ is not the only covariant quantity that can be connected, within a coordinate system, to the familiar comoving curvature perturbation of the linear theory. This is also the case for the projected gradient of the spatial scalar curvature (denoted by C_a in, e.g., [15]), although there does not seem to be a simple relation at the non-perturbative level between $\mathcal{R}_a$ and C_a .

Scalar field – It is also straightforward to extend our non-perturbative approach to a scalar field. Let us consider a scalar field ϕ with a potential $V(\phi)$. Its energy-momentum tensor, $T_{ab} = \partial_a\phi\partial_b\phi - \frac{1}{2}g_{ab}(\partial_c\phi\partial^c\phi + 2V)$, can be rewritten in the perfect fluid form by defining the four-velocity u^a as the unit vector orthogonal to hypersurfaces of constant ϕ [16],

$$u_a \equiv -\partial_a\phi/(-\partial_c\phi\partial^c\phi)^{1/2} = -\partial_a\phi/\dot{\phi}, \quad (16)$$

where the second equality (with $\dot{\phi} = u^a\partial_a\phi$) is a consequence of the definition of u^a , and the energy density and pressure as $\rho = \dot{\phi}^2/2 + V$ and $P = \dot{\phi}^2/2 - V$.

For a scalar field, the comoving curvature perturbation is particularly convenient because the $\phi = \text{const}$ hypersurfaces are orthogonal to u^a defined in (16). Since $D_a\phi = 0$, one sees that $\mathcal{R}_a = -D_a\alpha$ can be reexpressed as

$$\mathcal{R}_a = -\partial_a\alpha + \frac{\dot{\alpha}}{\dot{\phi}}\partial_a\phi,$$

and one recognizes the comoving curvature perturbation of the linear theory. The generalization of the Mukhanov variable [3] for the quantization of the scalar field-gravity system in the non-linear theory is thus $v_a \equiv e^\alpha(\dot{\phi}/\dot{\alpha})\mathcal{R}_a$.

In a coordinate system, $\mathcal{R}_a$ can be easily expanded to first and second orders to make contact with previous results. As for ζ_a , one finds, at first and second orders, expressions of the form

$$\mathcal{R}_i^{(1)} = \partial_i\mathcal{R}_1, \quad \mathcal{R}_i^{(2)} = \partial_i\mathcal{R}_2 + \frac{2}{\phi'}\delta\phi_1\partial_i\mathcal{R}'_1,$$

where the quantities $\mathcal{R}_1$ and $\mathcal{R}_2$ can be deduced from ζ_1 and ζ_2 , given in (8) and (9), by changing the overall sign and replacing ρ by ϕ . When first order vector and tensor modes can be neglected, up to small scale terms and an overall sign, $\mathcal{R} = \mathcal{R}_1 + \frac{1}{2}\mathcal{R}_2$ coincides with the variables used in [17] (see also [18]) either in the uniform field gauge, which can be obtained by setting $\delta\phi = 0$, or the uniform curvature gauge, obtained by setting $\delta\alpha = 0$. On writing α in terms of the metric components, and taking the large scale limit, one can also show that $\mathcal{R}$

can be simply related to the conserved quantities defined in [5, 7].

For a scalar field ($\omega_{ab} = 0$), if σ_{ab} can be neglected, one can also generalize the well known first order property that the nonadiabatic term Γ_a is negligible in the large scale limit. This result, which extends what is known up to second order [7], is obtained by noting that $D_a V = 0$ (since $D_a \phi = 0$), which implies $\Gamma_a = (1 - c_s^2)X_a \simeq 0$. In this case, for a *single scalar field*, the right hand side of Eq. (15) is negligible and $\mathcal{R}_a$ is conserved, on super-Hubble scales, to all orders in perturbation theory. If more fields are present, then $\Gamma_a \simeq -2D_a V \neq 0$ and the non-linear perturbation $\mathcal{R}_a$, and thus the non-Gaussianity, can evolve on large scales during inflation.

Conclusion – In this Letter, we have defined, respectively in (5) and (12), non-perturbative generalizations of the so called uniform energy density curvature perturbation and comoving curvature perturbation, using the integrated expansion α , which represents the number of e-folds of the local expansion measured by an observer comoving with the fluid.

Although our perturbed quantities are similar to those advocated in recent works based on the long wavelength approximation [8, 9, 10], their originality is that they are constructed geometrically, independently of any coordinate system. Furthermore, we have been able to derive evolution equations governing these quantities, which are *fully non-linear* and *exact at all scales*. In particular, Eq. (6), obtained directly from the local conservation of the energy-momentum tensor and independently of Einstein's equations, is valid in any metric theory of gravity, including scalar-tensor theories or induced four-dimensional gravity in brane-world scenarios. Thus, our relations generalize the familiar conservation laws of the linear theory for adiabatic perturbations.

Another advantage of our formulation is that it is rather easy to relate to the coordinate-based perturbative formulation. Interestingly, our extremely simple non-linear expressions, like (5), automatically contain the seemingly complicated second-order structure that has been obtained in the coordinate-based approach, which we can easily rederive and extend to higher orders.

Finally, let us stress that our formulation, since it keeps track of all small scale terms, may simplify the study of non-linear perturbations even inside the Hubble radius. This would be very useful, in particular to compute the

non-Gaussianity from inflation [17] or to study the back-reaction of small scale modes on larger scales.

We thus believe that this approach might represent a significant contribution to the study of primordial cosmological perturbations beyond the linear order.

Acknowledgments: We thank Marco Bruni for crucial comments on a preliminary version of this work and Misao Sasaki for useful discussions. F.V. thanks the Institut d'Astrophysique de Paris for kind hospitality.

-
- [1] J. M. Bardeen, Phys. Rev. D **22**, 1882 (1980).
 - [2] H. Kodama and M. Sasaki, Prog. Theor. Phys. Suppl. **78**, 1 (1984).
 - [3] V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rept. **215**, 203 (1992).
 - [4] V. Acquaviva, N. Bartolo, S. Matarrese and A. Riotto, Nucl. Phys. B **667**, 119 (2003); N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, Phys. Rept. **402**, 103 (2004).
 - [5] H. Noh and J. c. Hwang, Phys. Rev. D **69**, 104011 (2004).
 - [6] K. A. Malik and D. Wands, Class. Quant. Grav. **21**, L65 (2004).
 - [7] F. Vernizzi, Phys. Rev. D **71**, 061301 (2005).
 - [8] G. I. Rigopoulos and E. P. S. Shellard, Phys. Rev. D **68** 123518 (2003); G. I. Rigopoulos and E. P. S. Shellard, arXiv:astro-ph/0405185.
 - [9] E. W. Kolb, S. Matarrese, A. Notari and A. Riotto, arXiv:astro-ph/0410541.
 - [10] D. H. Lyth and D. Wands, Phys. Rev. D **68**, 103515 (2003); D. H. Lyth, K. A. Malik and M. Sasaki, JCAP **0505**, 004 (2005).
 - [11] G. F. R. Ellis and M. Bruni, Phys. Rev. D **40** 1804 (1989).
 - [12] M. Sasaki and E. D. Stewart, Prog. Theor. Phys. **95**, 71 (1996).
 - [13] D. Wands, K. A. Malik, D. H. Lyth and A. R. Liddle, Phys. Rev. D **62** 043527 (2000).
 - [14] G. F. R. Ellis, Relativistic Cosmology, in *General Relativity and Cosmology*, proceedings of the XLVII Enrico Fermi Summer School, edited by R. K. Sachs (Academic, New York, 1971).
 - [15] M. Bruni, P. K. S. Dunsby and G. F. R. Ellis, Astrophys. J. **395**, 34 (1992).
 - [16] M. Bruni, G. F. R. Ellis and P. K. S. Dunsby, Class. Quant. Grav. **9**, 921 (1992).
 - [17] J. Maldacena, JHEP **0305**, 013 (2003).
 - [18] D. S. Salopek and J. R. Bond, Phys. Rev. D **42**, 3936 (1990).